


Scorecard for Accountability and Action



Malaria is present in the three northern provinces of South Africa bordering Mozambique and Eswatini. The annual reported number of malaria cases in 2018 was 18,638 with 69 deaths.

Metrics

Commodities Financed and Financial Control	
IRS financing 2020 (% of at-risk population)	100
Public sector RDT financing 2020 projection (% of need)	100
Public sector ACT financing 2020 projection (% of need)	100
World Bank rating on public sector management and institutions 2018 (CPIA Cluster D)	
Insecticide Resistance Monitoring, Implementation and Impact	
Insecticide classes with mosquito resistance in representative sentinel sites confirmed since 2010	2
Insecticide resistance monitored since 2015 and data reported to WHO	
National Insecticide Resistance Monitoring and Management Plan	
Country Reporting Launch of Zero Malaria Starts with Me Campaign	
Scale of Implementation of ICCM (2017)	
Operational LLIN/IRS coverage (% of at risk population)	80
On track to reduce case incidence by $\geq 40\%$ by 2020 (vs 2015)	
On track to reduce case mortality by $\geq 40\%$ by 2020 (vs 2015)	
Tracer Indicators for Maternal and Child Health and NTDs	
Mass Treatment Coverage for Neglected Tropical Disease (NTD index, %)(2018)	2
Estimated % of Total Population living with HIV who have access to antiretroviral therapy (2018)	62
Estimated % of children (0–14 years old) living with HIV who have access to antiretroviral therapy (2018)	63
% deliveries assisted by skilled birth attendant	97
Postnatal care (within 48 hrs)	84
Exclusive breastfeeding (% children < 6 months)	32
Vitamin A Coverage 2018 (2 doses)	50
DPT3 coverage 2018 (vaccination among 0-11 month olds)	74

Key

	Target achieved or on track
	Progress but more effort required
	Not on track
	No data
	Not applicable

Malaria

Sustaining Essential Health Services During the COVID-19 Pandemic

The COVID-19 pandemic is putting an incredible strain on health systems across Africa. Health systems are required to maintain routine health services for other illnesses even as they handle the additional burden of COVID-19. In order to prevent widespread morbidity and mortality, it is of vital importance that we work to sustain the delivery of essential life-saving interventions during this difficult time including for Reproductive, Maternal, Newborn, Child and Adolescent health including malaria.

WHO underlines the critical importance of sustaining efforts to prevent, detect and treat malaria during the COVID-19 pandemic. It is of vital importance to ensure the continuity of malaria prevention and treatment services including distribution of insecticide-treated nets and indoor residual spraying. Any intervention must consider the importance of both lowering malaria-related mortality and ensuring the safety of communities and health workers given the ease of transmission of COVID-19.

For South Africa, it will be of vital importance that the indoor residual spraying (IRS) campaign planned for the fourth quarter of 2020 goes ahead and that the insecticides for the campaign are ordered on time, whilst taking into account physical distancing, in accordance with the recent guidance and recommendations from WHO. Without this IRS, coupled with the need to sustain essential health services including malaria case management, there could be an increase in malaria cases and deaths. Under the worst-case scenario, in which there is a 75% reduction in access to effective antimalarial medicines, WHO estimate that there could be a 20.6% increase in malaria cases, and a 135% increase in malaria deaths in South Africa. This scenario would represent a complete reversal in the substantial progress in malaria mortality reductions seen over the last 2 decades.

Progress

South Africa secured sufficient resources for the insecticides required for IRS in 2020 and enough RDTs and ACTs. WHO has identified South Africa as being a country with the potential to eliminate local transmission of malaria by 2020. The country has finalised the insecticide resistance monitoring and management plan. South Africa has significantly enhanced the tracking and accountability mechanisms for malaria with the development of the Malaria Elimination Scorecard.

Impact

The annual reported number of malaria cases in 2018 was 18,638 with 69 deaths.

Key Challenges

- Risk of re-introduction of malaria from neighbouring countries.
- There is a need to strengthen cross border collaboration with neighbouring countries.
- Sustaining the delivery of essential life-saving interventions during the COVID-19 pandemic including for Reproductive, Maternal, Newborn, Adolescent and Child health including malaria.

Previous Key Recommended Action

Objective	Action Item	Suggested completion timeframe	Progress	Comments - key activities/accomplishments since last quarterly report
Vector Control	Ensure the IRS programme is fully implemented before the beginning of the malaria season	Q4 2019		South Africa procured sufficient insecticide for the implementation of IRS in 2019, but spraying was delayed due to operational reasons and only completed at the end of March

New Key Recommended Action

Objective	Action Item	Suggested completion timeframe
Impact	Ensure that malaria services including case management and IRS are sustained and implemented whilst using COVID-19 sensitive guidelines during the pandemic	Q4 2020

RMNCAH and NTDs

Progress

The country has made significant progress in scaling-up the tracer RMNCAH interventions of skilled birth attendants and postnatal care.

Progress in addressing Neglected Tropical Diseases (NTDs) in South Africa is shown using a composite index calculated from preventive chemotherapy coverage achieved for schistosomiasis and soil transmitted helminths. Preventive chemotherapy coverage in South Africa is below WHO target for soil transmitted helminths (54%) and preventive chemotherapy for schistosomiasis was not implemented in 2018 (0%). Overall, the NTD preventive chemotherapy coverage index for South Africa in 2018 is very low (2), which is the same as in 2017 index value (2).





Previous Recommended Actions

Objective	Action Item	Suggested completion timeframe	Progress	Comments - key activities/accomplishments since last quarterly report
NTDs	Identify the reasons for the decrease in Preventive Chemotherapy (PC) coverage of Soil Transmitted Helminths (STH) and the continued low coverage of PC for Schistosomiasis, and work to enhance coverage to reach the WHO targets	Q4 2019		The programme has carried out the mapping of Schistosomiasis distribution in 7 provinces with only 2 outstanding. This will support the development of a national strategy to treat everyone everywhere in need of preventive chemotherapy for NTDs
NTDs	Implement preventive chemotherapy (PC) for schistosomiasis along with soil transmitted Helminths (STH) and work to increase PC for STH to reach WHO target.	Q2 2020		Deliverable not yet due

New Key Recommended Action

Objective	Action Item	Suggested completion timeframe
RMNCAH ¹ : Impact	Ensure that essential RMNCAH services are sustained and implemented whilst using COVID-19 sensitive guidelines during the pandemic	Q4 2020

Key

	Action achieved
	Some progress
	No progress
	Deliverable not yet due

¹ RMNCAH metrics, recommended actions and response tracked through WHO